Two thwest Regions. Summary of Rej Two Hold Till Harings			erbera IDPs, Post <i>Deyr</i> 2010-11				
	Hargeisa IDPs N=651		Burao IDPs N=652		Berbera IDPs N= 542		
Indicator	Results	Outcome	Results	Outcome	Results	Outcome	
Child Nutrition Status							
Global Acute Malnutrition (WHZ<-2 or oedema)	10.8 (8.9-13.0)	Serious	12.1 (8.8-15.4)	Serious	14.2 (10.9-18.3)	Serious	
Boys	12.0 (9.7-14.8)	_	14.1 (9.2-15.8)	_	16.3 (12.2-21.5)		
Girls	9.3 (6.0-14.1)		9.8 (6.7-13.0)		12.3 (7.7-19.0)		
Severe Acute Malnutrition (WHZ<-3 or oedema)	1.5 (0.8-2.8)	Acceptable	1.7 (0.8-3.4)	Acceptable	2.2 (1.2-4.1)	Acceptable	
Boys	2.0 (1.0-3.9)		2.0 (0.5-3.6)		1.6 (0.6-3.9)		
Girls	1.0 (0.2-4.4)		1.3 (0.1-2.6)		2.8 (1.3-5.8)		
Mean Weight for height Z score	-0.57	Alert	-0.65	Alert	-0.80	Serious	
Oedema		Alert		Alert	0	Alout	
	0.2 (0.0-0.5)		0.5 (0.0-1.2)			Alert	
Acute malnutrition by MUAC (<12.5 cm or oedema)	4.6 (2.9-7.3)	Acceptable	4.4 (2.9-6.0)	Acceptable	9.2 (5.1-13.3)	Serious	
Boys	4.6 (2.7-7.6)		3.7 (1.7-5.8)		8.0 (4.3-11.7)		
Girls	4.6 (2.1-9.8)		5.2 (3.1-7.4)		10.4 (5.0-15.8)		
Acute malnutrition by MUAC (<11.5 cm or oedema)	1.5 (0.7-3.6)	Serious	0.8	Alert	3.3	Critical	
Global Acute Malnutrition (WHZ<-2 or oedema) NCHS	9.7	Alert	11.0	Serious	14.0 (10.6-18.3)	Serious	
Severe Acute Malnutrition (WHZ <-2 or oedema) NCHS	0.8	Alert	0.6	Alert	1.1 (0.5-2.4)	Alert	
Stunting (HAZ<-2)	10.4 (7.1-15.1)	Serious	4.6 (2.1-7.1)	Alert	6.1 (4.0-8.1)	Alert	
Boys	13.8 (9.5-19.5)		6.3 (2.8-9.9)		8.6 (5.3-11.8)		
Girls	6.6 (3.8-11.2)		2.6 (0.6-4.6)		3.9 (1.4-6.3)		
Underweight (WAZ<-2)	9.4 (6.4-13.6)	Alert	9.2 (6.5-11.9)	Alert	12.7 (9.8-15.7)	Serious	
		THOIT		THOIT		Serious	
Boys	12.4 (7.9-18.9)		12.4 (8.2-16.6)		17.1 (13.8-21.2)		
Girls	6.0 (3.5-10.0)		5.6 (3.5-7.7)		8.8 (4.6-13.0)		
HIS Nutrition Trends(Jan-Jun '10)	Low (<10%) and	Serious	Low (<10%) and	Serious	Low (<10%) and	Serious	
	stable trend of		stable trend of		stable trend of		
	acutely		acutely		acutely		
	malnourished		malnourished		malnourished		
	children		children		children		
Proportion of acutely malnourished children in SFs	3.6	Very Critical	9.1	Very Critical	0.8	Very Critic	
Child Morbidity & Immunization							
Disease trends (seasonally adjusted)	Morbidity- 34.4	Critical	Morbidity- 32.7	Critical	Morbidity- 35.6	Critical	
Morbidity refers to the proportion of children reported to be ill in the 2	Diarrhoea – 16.9	OTTION.	Diarrhoea – 16.6	Critical	Diarrhoea – 21.8	Critical	
	Pneumonia- 11.7		Pneumonia- 13.0		Pneumonia- 11.3		
weeks prior to the survey							
Immunization Status	Vitamin A – 76.3	Critical	Vitamin A – 69.5	Critical	Vitamin A – 75.8	Critical	
	Measles – 73.9		Measles – 69.6		Measles – 72.7		
Infant and Young Child Feeding	N= 236		N=235		N=275		
Proportion still breastfeeding	40.3	Critical	39.6	Critical	38.2	Critical	
Proportion meeting recommended feeding frequencies	35.2	Critical	48.1	Critical	28.8	Serious	
Proportion who reported to have consumed >4 food groups	14	Very Critical	3.4	Critical	3.9	Critical	
Death Rates		, and the second					
Crude deaths per 10,000 per day 9retrospective for 90 days)	0.36 (0.21, 0.62)	Aggantable	0.47 (0.29-0.75)	Accentable	0.12 (0.05.0.22)	Aggentable	
Crude deaths per 10,000 per day steriospective for 90 days)	0.36 (0.21- 0.62)	Acceptable	0.47 (0.29-0.73)	Acceptable	0.12 (0.05-0.33)	Acceptable	
Under five deaths per 10,000 per day retrospective for 90 days)	0.95 (0.44-2.04)	Acceptable	1.36 (0.68-2.69)	Acceptable	0.33 (0.08-1.35)	Acceptable	
Women Nutrition and Immunization Status	N=40')	N=406		N=3	76	
Proportion of acutely malnourished non pregnant/lactating women	0	Acceptable	0	Acceptable	0.5	Acceptable	
(MUAC <18.5 cm)							
Proportion of acutely malnourished pregnant and lactating women	15.4	Serious	20.6	Serious	16.1	Serious	
(MUAC<23.0)	15.1	Serious	20.0	Serious	10.1	Serious	
Proportion of Women who received Tetanus immunization		Serious		Serious		Serious	
No dose	16.7		26.1		14.4		
One dose	15.5		18.5		14.7		
Two doses	28.0		26.8		35.1		
Three doses	39.8		28.6		35.8		
Public Health Indicators	N= 42	3	N=42	7	N=3	79	
Household with access to sanitation facilities	77.1	Critical	70.0	Critical	20.3	Critical	
rousehold with access to sailtation facilities	//.1	Criucai	70.0	Critical	20.3	Crucal	
Household with access to safe water	93.9	Serious	88.5	Serious	96.9	Alert	
Food Security							
Proportion who reported to have consumed <4 food groups	12.3	Alert	21.5	Serious	12.9	Alert	
Household's Main Food Source							
Own production	-		-		_		
Purchase	97.5		96.2		98.9		
	11.3		70.2		70.7		
Food security phase	BFI	Serious	BFI	Serious	BFI	Serious	

Overall Risk to Deterioration	Unstable	Serious	Unstable	Serious	Unstable	Serious
Overall Situation Analysis	Seriou	S	Seriou	S	Serio	ous