

# Somalia 2020 Post *Deyr* Food Security and Nutrition Outcomes and Projections

A Virtual Briefing for All Stakeholders

4 February 2021

Funding for the 2020 Post *Deyr* Assessments and subsequent IPC analyses was provided by:



**USAID**  
FROM THE AMERICAN PEOPLE



**FSNAU Donors**



This project is co-funded  
by the European Union



Schweizerische Eidgenossenschaft  
Confédération suisse  
Confederazione Svizzera  
Confederaziun svizra

Swiss Agency for Development  
and Cooperation SDC


# 2020 Post-Deyr Assessment, Analysis and Vetting Process

Planning, assessment, analysis and vetting of the results were conducted in collaboration with government, UN agencies, local and international NGOs and technical partners.


- Briefing of partners on 2020 Post Deyr assessment plan and survey protocol – **Oct/Nov 2020**
- Regional Planning Workshop/Training for rural food security assessment: Hargeisa, Garowe, Galkacyo, Dhusamareb, Beletweyne, Mogadishu, Baidoa, Dollow and Kismayo: **Dec 2020**
- Fieldwork (data collection): **Nov-Dec 2020**
- Regional Analyses Workshops: **6-10 January 2021**
- IPC Analyses Workshops and Technical Vetting: **11-19 Jan**
- Technical Briefing for Government (virtual): **1 Feb**
- Briefing for UN Heads of Humanitarian Agencies (virtual): **2 Feb**
- Briefing for Senior Government Officials: (virtual): **3 Feb**
- Final Dissemination to All Stakeholders (virtual): **4 Feb**
- **Participation in the 2020 Post Deyr IPC Analysis:**
- Total number of participants: **103 participants**
  - Government institutions: **33 participants**
    - Federal Government of Somalia
    - Galmudug
    - Hirshabelle
    - Southwest
    - Jubaland
    - Puntland
    - Somaliland
  - NGOs/INGOs: **25 participants**
  - Local Universities (Puntland State University): **2 participants**
  - Technical partners (FEWS NET, IPC GSU and): **5 participants**
  - UN (FAO/FSNAU, WFP and UNICEF): **35 participants**
  - Food Security and Nutrition Clusters - **3**

# 2020 Deyr Season Rainfall and Temperature Forecast

Oct-Dec 2020 Rainfall Outlook  
(Probabilistic Forecast)




Oct-Dec 2020 Temperature Outlook  
(Probabilistic Forecast)




- Forecast issued by the Greater Horn of Africa Climate Outlook Forum (GHACOF56) through IGAD/ICPAC in late August 2020 indicated a greater likelihood of below normal to normal *Deyr* season (Oct-Dec 2020) rainfall in most parts of Somalia and average to below average rainfall in northeast regions.
- Below average Xays (Dec-Jan) rainfall were also considered likely in northwestern Somalia.
- Warmer than usual temperatures were expected over most of the Greater Horn of Africa (GHA) region, including Somalia.

# 2020 Deyr Season Rainfall Performance

Cumulative Seasonal Rainfall in MM (CHIRPS), Oct-Dec 2020



Cumulative Seasonal Rainfall Deviation from Average in MM (CHIRPS), Oct-Dec 2020




- The 2020 Deyr season rainfall was characterized by delays and erratic distribution Overall, the 2020 Deyr season rainfall was below average in most parts of the country with adverse impact on replenishment of pasture and water resources and on Deyr season crop production.
- Mixed 2020 Deyr onset; delays in most of North and large parts of the South (Gedo, Shabelle, and Juba regions);

- Ineffective rainfall establishment in the North, erratic distribution across many areas.
- Cyclone Gati with torrential rains which hit coastal areas of Bari (late November) caused significant damages although it also brought favorable rainfall to coastal and adjacent pastoral areas of Sanaag, Woqooyi Galbeed and Awdal.
- Rainfall totals in most northern and southern areas below average.
- Average to above average totals in many areas of central regions, parts of Bay, Bakool and Hiiraan and in Cyclone GATI affected areas in the northeast and adjacent parts of northwest,
- Floods continued in Hiiraan, Shabelle and Juba regions from late October to early November

# 2021 Gu Season Rainfall Forecast

## March-May 2021 Rainfall Outlook



- In the Horn of Africa (GHA) region, La Niña is usually associated with drought conditions and El Niño with flooding. GHA climate is also influenced by the Indian Ocean Dipole (IOD).
- Latest forecasts from CPC/NCEP/NWS/IRI indicates that La Niña is expected to continue during *Jilaal* (~95% chance between January and March 2021), with a potential transition to ENSO-neutral (i.e. neither La Niña, nor El Niño) during the forthcoming Gu (55% between April and June 2021).
- Based on assumptions of waning La Niña conditions, warm western Pacific Ocean gradients, and a neutral IOD, there is greater likelihood of below-average to average April to June 2021 Gu season rainfall in central and southern Somalia and below average rainfall in the northern regions.

# Market Prices

## July-December 2020


- Somali/Somaliland Shilling was stable over the past six months, except in Puntland where the local Somali Shilling continued to depreciate against the US Dollar.
- Prices of imported food commodities were generally stable, except in in northeast (Puntland) where prices increased sharply due to depreciation of the local currency.
- Cereal food prices (maize and sorghum) were generally stable and close to the five-year average apart from the northeast region that has been affected by inflationary pressures caused by depreciation of the local currency.
- Significant increase in the cost of the Minimum Expenditure Basket (MEB) was observed in the Puntland regions.
- Livestock exports declined sharply between August to December 2020 compared to average.




## January-June 2021 Outlook

- The SOSH/SLSH exchange rate against USD is likely to be stable in most markets, partly boosted by expected increases in livestock export in the lead up to and during Ramadan and Hajj (March-May). However, some exchange rate fluctuations is expected in northeast markets.
- A below average 2020 Deyr harvest and expected below average 2021 Gu season rainfall will likely tighten domestic cereal supply and put upward pressure on prices starting in April.
- Sorghum and maize imports from Ethiopia will supplement supplies to adjacent markets in northern, central and southern regions.
- Food import (rice, flour and sugar) prices are expected to be stable through mid-2021 from promising global export availabilities, sluggish demand and low oil price.
- Livestock prices will likely follow seasonally trend through June from improved livestock condition and productivity as well as increased exports. Prices are expected to be close or above the average in most markets with regional differences.

# Impact on Agriculture (Maize and Sorghum Production)




- In southern Somalia, the 2020 *Deyr* season cereal production is estimated at **78 600 tons**, including **4 100 tons** of off-season harvest expected in March 2021. The 2020 *Deyr* harvest is **20** percent lower than the long-term average for 1995-2019.
- Main factors for the reported production decline include: erratic rainfall, successive and severe flooding, Desert Locust and insecurity/conflict.



- In northwest regions, the 2020 *Gu/Karan* cereal production was previously estimated at **22 500 MT** (September 2020), based on mid-season crop assessment.
- However, based on crop harvest assessments conducted in November, the revised production estimate is **17 100 MT**. This is **58** percent lower than the average for 2010-2019, mainly due to erratic rainfall, Desert Locust and stalk borer infestation on both sorghum and maize crops.

# Impact on Pasture and Water Availability



- Carryover water and pasture from the 2020 *Gu* season supported livestock through late October. Despite Desert Locust damage in northern and central and parts of south regions, average to near average 2020 *Deyr* rainfall in November have moderated the impact and replenished pasture and browse in south/central and parts of north.
- In late November, tropical cyclone GATI was experienced in northern regions mainly, Hafuun/Iskushuban, Alula, Bederbeyla, Bossasso, Qandala, Erigavo and Laasqoray districts. Although the cyclone caused significant damage, it also improved pasture and water in the affected areas.
- Harsh *Jiilaal* (January-March 2021) is expected particularly in rain deficit areas of northern and central regions, leading to water scarcity, increased water trucking and early and atypical livestock migration and increased abortion among small ruminants in some regions.


## Impact on Livestock Production and Productivity

---

- Conception during the 2020 Deyr was medium among small ruminants. Among large ruminants, conception was low to medium during the 2020 Deyr due to long gestation periods (most of them conceived 2020 *Gu* seasons or have given birth during the 2020 Deyr). Livestock births during the 2020 Deyr were also medium for small ruminants.
- Milk availability is average to above average in southern Somalia, except in Gedo where milk availability is low due to less favorable pasture and browse conditions.
- Although milk production has improved in northern and central regions, availability remains low due to (1) low ownership of milking/lactating animals as herd sizes have not yet fully recovered from the extended cumulative impact of previous droughts (2) most livestock have migrated to distant grazing areas in search of water and pasture.
- Milk availability is expected to decline through mid-2021 due to harsh conditions during *Jilaal* and a likely below average 2021 *Gu* (April-June) season rainfall, particularly in northern and central regions.
- Livestock holding among poor pastoral households continued to increase during the 2020 Deyr season. Further increases are expected towards mid-2021 year due to anticipated medium births between now and mid-2021 in southern regions.
- However, livestock holding among poor pastoral households will still remain below baseline in central and parts of northern Somalia by mid-2021 but reach near baseline or above baseline levels in southern Somalia.

# Desert Locust Infestation and Outlook


Aug 2020




Sep 2020



Oct 2020




Nov 2020




- Desert locust which has mostly affected northern and central regions since late 2019, it has expanded to southern Somalia since October 2020.
- The southward movement and breeding of Desert Locust coincided with both the planting and harvesting periods. Despite ongoing control operations which are reducing impacts.


Dec 2020



4-10 Jan 2021



Forecast (Jan-Mar 2021)





- Desert Locust has caused significant damage to 2020 Deyr season cereal crop cultivation and harvest in central and southern regions.


- Latest forecasts indicate that Desert Locust continues to pose a very high risk to rural livelihoods across Somalia in 2021, potentially exacerbating the impact of a below average 2021 *Gu*.

Source of Maps and Forecast: FAO Desert Locust Watch

# Somalia Population Movement/Displacement and Impact, Jul-Dec 2020 (UNHCR/PRMN Data)




- Overall, nearly **661 000** persons were displaced between July and Dec 2020, mainly due to floods (**69%**), insecurity/conflict (**14%**) and lack of livelihoods (**14%**).




- Most of the displacements occurred between July and Sept and November, mainly driven by recurrent floods.
- Most of the population displacements occurred in Lower Shabelle and Middle Shabelle and Banadir regions.
- Flooding and insecurity/conflict related displacements have contributed to lower crop production in Hiran, Middle and Lower Shabelle regions.
- Armed and political conflict, drought and flood related displacements are expected to continue through mid-2021, further exacerbating food insecurity across many parts of Somalia.

# Remittance Receipts by Households, Aug/Sep-Oct/Nov 2020


**% of HHs who received cash remittances (Rural)**




**% of HHs who received cash remittances (Urban and IDPs)**




**Average amount received in USD (Rural)**




**Average amount received in USD (Urban and IDPs)**



**If remittance is received how it compares with typical amounts (Rural)**



**If remittance is received how it compares with typical amounts (Urban and IDP)**




- Up to 22% of urban, 12% of rural and 6% IDPs received remittances between Aug/Sep and Oct/Nov 2020.


- Amount recipients, average amounts are significant in many areas

- A majority of recipients (>50%) reported significant declines in remittances

# 2020 Post Deyr Nutrition Situation: GAM and SAM



At the national level, the median prevalence of Global Acute Malnutrition (GAM) has remained Serious (10–14.9%) over the past three seasons (11.8% in 2020 Deyr; (11.8% in 2020 *Gu* and 13.1% in 2019 *Deyr*).




	GAM
Extremely Critical	≥30%
Critical	15.0-29.9%
Serious	10.0-14.9%
Alert	5.0-9.9%
Acceptable	<5%


	SAM
Extremely Critical	>5.6%
Critical	4.0-5.6%
Serious	2.5-3.9%
Alert	1.0-2/4%
Acceptable	<1%

# Morbidity, Vit A Supplementation and Measles Vaccination among Children Under-Five


## Morbidity (%)



## Vitamin A Supplementation (%)



## Measles Vaccination (%)



# Nutrition Situation in Rural Livelihoods

Population Group	Acute Malnutrition Prevalence				Per 10 000 per Day		Morbidity Among Children Under- Five (%)	Children Under-Five		Household	
	Children Under-Five		Women Aged 15-49		Crude Death Rate (CDR)	Under-Five Death Rate (U5DR)		Coverage (%)		Access to	
	GAM (%)	SAM (%)	MUAC < 23 CM (%)	MUAC < 21 CM (%)				Vit A Suppl.	Measles Vaccination	Clean Water (%)	Sanitation (%)
Guban P	5.9	0.6	10.7	3.3	0.30	0.15	22.4	29.8	29.3	66.0	53.0
West Golis P	7.8	0.7	6.9	0.5	0.31	0	18.3	64.4	56.8	68.0	53.0
Northwest AP	5.9	0.5	0.9	0	0.16	0	25	37.9	36.5	15.0	38.0
NIP of NW	11.2	2.0	15.2	7.1	0.62	0.35	7.8	73.1	72.6	33.0	84.0
Hawd P-NW	6.6	1.3	17.7	10	0.35	0	6.2	90.7	90.5	32.0	90.0
East Golis P	11.8	1.9	29.9	12.7	0.34	0.27	42.7	67.9	71.1	46.0	46.0
NIP of NE	7.7	1.0	33.3	4.1	0.05	0	17.7	51.4	50.9	47.0	100.0
Hawd P of Central	14.9	3.5	21.0	13.3	0.27	0.57	27.7	59.7	57.2	27.0	67.0
Coastal Deeh P	5.1	0.0	7.1	1.6	0.06	0.21	24.6	47.5	50	71.0	10.0
Addun P	10.4	1.5	12.0	2.6	0.15	0.23	21.1	44.3	44.8	24.0	71.0
Shabelle Riv	15.4	2.5	16.1	13.1	0.76	1.51	14.6	27.7	9.0	22.0	55.0
Shabelle API	14.9	2.9	19.3	2.1	0.47	1.07	16.8	27.3	16.1	45.0	44.0
Bay AP	14.4	2.4	19.5	4.6	0.30	0.4	34.9	21.6	23.4	4.0	6.0
Bakool P (ACF)	20.9	4.1	25.0	8.7	0.04	0	24.7	40.9	49.5	19.0	42.9
N Gedo P	12.5	1.7	17.9	8.1	0.31	0.48	26.6	59.4	54.3	10.0	24.0
N Gedo Riv	14.8	1.9	23.6	3.4	0.36	0.81	31.1	78.9	70.0	39.0	66.0
<b>Median</b>	<b>11.5</b>	<b>1.8</b>	<b>17.8</b>	<b>4.4</b>	<b>0.3</b>	<b>0.3</b>	<b>23.5</b>	<b>49.5</b>	<b>50.5</b>	<b>32.5</b>	<b>53.0</b>

- There has been some improvement in the overall nutrition situation among rural populations compare to 2019 *Deyr* (13.8 % to 11.5%). Critical (GAM ≥ 15%) was observed only in two out of 16 population group;s surveyed (Shabelle riverine and Bakool Pastoral).
- Morbidity prevalence were high (≥ 20%) for most (10 out of 16) of rural population groups.
- Crude Death Rate (CDR) and Under-Five Death Rate (U5DR) were low across most rural livelihoods. Exceptions were Shabelle riverine which recorded Serious level of CDR (0.5-1/10 000/day) and U5DR (1 to1.9/10 00 /day. Serious CDR was also observed in Northern inland Pastoral and Shabelle Agropastoral.
- Measles vaccination, Vitamin A supplementation and household access to clean water and sanitation remain low in many rural livelihoods.

# Nutrition Situation Among Urban Populations

Population Group	Acute Malnutrition Prevalence				Per 10 000 per Day		Morbidity Among Children Under- Five (%)	Children Under-Five		Household	
	Children Under-Five		Women Aged 15-49		Crude Death Rate (CDR)	Under-Five Death Rate (U5DR)		Coverage (%)		Access to	
	GAM (%)	SAM (%)	MUAC < 23 CM (%)	MUAC < 21 CM (%)				Vit A Suppl.	Measles Vaccination	Clean Water (%)	Sanitation (%)
Hargeisa Urban	6.4	1.1	1.4	0.9	0.16	0	23.5	66.1	55.4	100.0	94.0
Burao Urban	4.3	0.5	2.2	0	0.43	0	4.3	83.9	83.1	100.0	100.0
Bosasso Urban	7.5	1.3	6.7	3.1	0.04	0.18	17.7	73.8	72.2	94.0	98.0
Garowe Urban	5.2	0.3	2.3	0.9	0.34	0.16	25.2	82.7	80.6	62.0	99.0
Galkacyo Urban	11.8	1.4	7.9	2.4	0.18	0	24.0	58.7	58.4	95.0	100.0
Dhusamareb Urban	5.8	0.2	12.7	6.6	0.00	0	17.7	55.3	50.9	100.0	96.0
Mogadishu Urban	13.2	2.3	14.9	10.4	0.6	0.58	16.4	72.7	69.9	100.0	100.0
Huddur Urban	16.4	5.8	6.9	2.3	0.5	0.36	25.1	49.7	91.0	18.4	81.0
Baidoa Urban	9.0	1.2	19.5	2.8	0.24	0.32	33.6	68.9	60.6	96.0	99.0
Dolow Urban	12.4	0.8	13.4	3.7	0.29	0.66	26.9	84.9	80.5	98.0	100.0
Kismayu Urban	12.2	2.4	10.8	3.2	0.35	0.52	16.8	90.5	90.2	86.0	100.0
<b>Median</b>	<b>9.0</b>	<b>1.2</b>	<b>7.9</b>	<b>2.8</b>	<b>0.3</b>	<b>0.2</b>	<b>23.5</b>	<b>72.7</b>	<b>72.2</b>	<b>96.0</b>	<b>99.0</b>

- Morbidity among children was high (Over the past three seasons, there has been improvement in the overall nutrition situation among urban populations: 9.0% in 2020 Deyr compared to 10.0% in 2020 Gu and 11.5% in 2019 Deyr).
- A Critical GAM prevalence of 16.4% level was only reported among urban populations in Hudur.
- Morbidity prevalence is high ( $\geq 20\%$ ) in 6 out of 11 urban populations, with the highest prevalence reported among Baidoa urban (33.6%); Dollow urban (26.9%) and Garowe urban (25.2%).
- Crude Death Rate (CDR) and Under-Five Death Rate (U5DR) were low across most urban population groups, with the exceptions of Mogadishu urban and Huddur urban which had a Serious level of CDR.

# Nutrition Situation Among IDP Populations

Population Group	Acute Malnutrition Prevalence				Per 10 000 per Day		Morbidity Among Children Under- Five (%)	Children Under-Five		Household	
	Children Under-Five		Women Aged 15-49		Crude Death Rate (CDR)	Under-Five Death Rate (U5DR)		Coverage (%)		Access to	
	GAM (%)	SAM (%)	MUAC < 23 CM (%)	MUAC < 21 CM (%)				Vit A Suppl.	Measles Vaccination	Clean Water (%)	Sanitation (%)
Hargeisa IDPs	9.2	1.2	2.8	0.7	0.24	0.33	22.7	30.1	28.8	100.0	99.0
Burao IDPs	5.8	1.8	2.7	0.9	0.09	0	4.7	83.1	77.8	99.0	100.0
Bosasso IDPs	15.8	2.5	22.8	4.9	0.27	0.67	19.4	73.9	75.4	63.0	88.0
Garowe IDPs	15.8	1.9	2.1	0	0.68	1.04	28.3	85	83.3	96.0	67.0
Galkacyo IDPs	10.8	2.2	13.8	5	0.23	0.27	32.1	65.3	65.3	96.0	91.0
Mogadishu IDPs	16.7	3.7	12.1	4.9	0.83	1.46	22.7	61.2	62.7	99.0	81.0
Baidoa IDPs	11.9	2.1	26.5	5.4	0.3	0.1	41.7	63.5	54.4	64.0	93.0
Dolow IDPs	13.6	1.6	7.4	3.1	0.3	0.9	24	79	74.8	99.0	100.0
Kismayu IDPs	12.9	2.3	7.0	1.5	0.4	0.8	16.5	94.2	92.2	89.0	91.0
<b>Median</b>	<b>12.9</b>	<b>2.1</b>	<b>7.4</b>	<b>3.1</b>	<b>0.3</b>	<b>0.7</b>	<b>22.7</b>	<b>73.9</b>	<b>74.8</b>	<b>96.0</b>	<b>91.0</b>

- Overall , the median GAM (WHZ) prevalence among Internally Displaced Populations (IDPs) in 2020 Deyr is Serious (12.9%), reflecting a slight improvement but sustained Serious nutrition situation since late 2019 (13.7% 2019 Deyr and 13.1 % in 2020 Gu).
- Critical prevalence of acute malnutrition persisted in most of the IDPs with the exception of Hargeisa and Burao since Deyr 2019. These results underscore the underlying vulnerability of IDP populations.
- Morbidity prevalence were high ( $\geq 20\%$ ) among majority (6 out of 9) of IDP population groups.

- Crude Death Rate (CDR) and Under-Five Death Rate (U5DR) were low across most IDP population groups. Exceptions were IDPs in Garowe and Mogadishu which had Serious level of CDR and/or U5DR. Measles vaccination, Vitamin A supplementation are low in several of the IDP population groups. However, household access to clean water and sanitation is high among most of the displaced populations.

# Results of MUAC Assessments Among Children in Hard to Reach Areas

- Only three out of 9 MUAC screened areas are Classified as **Critical** , while 6 areas classified as **Serious**.

- Out of 9 surveyed areas, 5 recorded morbidity prevalence > 20%.

Population Groups	Acute Malnutrition Prevalence Among Children Under-Five		Morbidity Among Children Under-Five (%)
	MUAC < 12.5 CM (%)	MUAC < 11.5 CM (%)	
Beletwein District (Riverine)	7.2	1.0	14.3
Beletwein Urban	9.0	1.3	12.9
South Gedo Pastoral	8.4	1.3	15.1
South Gedo Riverine	6.9	1.5	20.8
South Gedo Agropastoral	5.9	1.2	15.8
Juba Cattle Pastoral	7.4	0.9	21.1
Juba Agropastoral	9.2	1.3	20.4
Juba Riverine	13.7	2.6	32.3
Elberde Pastoral (Bakool)	9.2	1.0	23.5

## Summary Results for Women of Child Bearing Age (15-49 years old)


- Acute malnutrition is also high among women of child bearing age (25-49 years old) in many areas.
- Out of the 44 population groups surveyed, six had Critical acute malnutrition prevalence (23.4-31.4%), 10 were Serious (GAM 16.8-23.3%) while 10 were Alert (GAM 10.5-16.7).

Acute Malnutrition Prevalence			Acute Malnutrition Prevalence		
Women Aged 15-49			Women Aged 15-49		
Population Group	MUAC < 23 CM (%)	MUAC < 21 CM (%)	Population Group	MUAC < 23 CM (%)	MUAC < 21 CM (%)
Guban Pastoral	10.7	3.3	Beletwein District (Riverine)	10.8	2.6
West Golis Pastoral	6.9	0.5	Beletwein Urban	17.1	2.2
Northwest Agropastoral	0.9	0	Shabelle Riverine	16.1	13.1
Hargeisa IDPs (W. Galbeed)	2.8	1	Shabelle Agropastoral	19.3	2.1
Hargeisa Urban (W. Galbeed)	1.4	0.9	Mogadishu Urban	14.9	10.4
Burao IDPs (Toghdeer)	2.7	0.9	Mogadishu IDPs	12.1	4.9
Burao Urban (Toghdeer)	2.2	0	Bay Agropastoral	19.5	4.6
Northern Inland Pastoral of NW	15.2	7.1	Baidoa IDPs (Bay)	26.5	5.4
Hawd Pastoral of Northwest	17.7	10	Huddur Urban	6.9	2.3
East Golis Pastoral	29.9	12.7	Baidoa Urban (Bay)	19.5	2.8
Bosasso IDPs (Bari)	22.8	4.9	Dolow IDPs (N Gedo)	7.4	3.1
Bosasso Urban (Bari)	6.7	3.1	Dolow Urban (N Gedo)	13.4	3.7
Northern Inland Pastoral of NE	33.3	4.1	North Gedo Pastoral	17.9	8.1
Hawd Pastoral of Central	21.0	13.3	North Gedo Riverine	23.6	3.4
Coastal Deeh Pastoral of NE	7.1	1.6	Kismayu Urban (L. Juba)	10.8	3.2
Garowe IDPs (Nugaal)	2.1	0	Kismayu IDPs (L. Juba)	7.0	1.5
Garowe Urban (Nugaal)	2.3	0.9	South Gedo Pastoral	18.9	5.8
Galkacyo IDPs (Mudug)	13.8	5	South Gedo Riverine	20.0	11.7
Galkacyo Urban (Mudug)	7.9	2.4	South Gedo Agropastoral	32.0	14.0
Dhusamareb Urban (Galgadud)	12.7	6.6	Juba Cattle Pastoral	4.9	0.7
Addun Pastoral	12.0	2.6	Juba Agropastoral	9.1	4.5
Elberde Pastor (Bakool)	23.7	2.2	Juba Riverine	6.4	4.1

# Nutrition Key Messages

- Acute malnutrition remains high in many parts of Somalia.
- Morbidity prevalence among children are high (above 20%) in a majority of the population groups surveyed, including East Golis Pastoral (42.7%), Baidoa IDPs, Bay (41.7 %), Bay Agropastoral (34.9%), Baidoa Urban (33.6%), Galkacyo IDPs, (32.1%) and North Gedo Riverine (31.1%).
- Vitamin A Supplementation and measles vaccination: coverage status of these essential health/nutrition services is lowest (<40%) in Shabelle Agro-pastoral and Riverine in Middle and Lower Shabelle regions, Bay Agro-pastoral, Northwest Agropastoral, Guban pastoral and Hargiesa IDP in Northwest regions.
- Projection for February-April 2021: Level of acute malnutrition is projected to worsen across Somalia with five population groups expected to deteriorate from Serious to Critical, three from Alert to Serious and one from Acceptable to Alert. Other areas will worsen but sustain current phases of acute malnutrition.
- Hotspot Areas- Populations with Critical level of acute malnutrition in the current or projected period include Riverine population in both Shabelle and Juba regions, Agropastoral in Bay region, Hiran region, Pastoral population in Elberde district and Hawd of Central and IDPs in Bosasso, Mogadishu, Garowe, Baidoa and Galkayo.
- The situation in the current hotspot areas and most of the IDP settlements calls for sustained humanitarian interventions in the form of integrated nutrition but also include nutrition sensitive programs with special focus on resilience and prevention
- Coverage of health and nutrition services (both treatment and prevention), including vitamin A supplementation and measles vaccination should be expanded, especially in areas where coverage is currently low.

# Nutrition Outcomes and Projections: 2020 Post *Gu* and 2020 Post *Deyr*



Total Burden (Sep 2020–Aug 2021): **849 900** acutely malnourished, including **143 400** who are likely to be severely malnourished


Total Burden (Jan 2021–Dec 2021): **838 900** acutely malnourished, including **143 200** who are likely to be severely malnourished

- Based on results of the 2020 Post Deyr assessment, an estimated **838 900** children under the age of five years (total acute malnutrition burden) face acute malnutrition between January and December 2021, including **143 200** likely to be severely malnourished. This reflects a slight improvement in the overall nutrition situation and outlook compared to 2019 *Deyr* and 2020 Post *Gu*.

## Key Food Security Related Findings

- The 2020 Deyr season rainfall was characterized by delays and erratic distribution. Overall, the 2020 Deyr season rainfall was below average in most parts of the country with adverse impact on replenishment of pasture and water resources and on Deyr season crop production. This in turn has adversely affected food security outcomes among farmers and pastoralists. Humanitarian assistance (food security and nutrition) and government support have prevented the worsening of food security and nutrition outcomes across many parts of Somalia.
- Poor pastoral households in northern and central Somalia continue to face moderate to large food consumption gaps as they have limited number of livestock to help them cope with the various shocks both during the current and projection periods.
- In agropastoral livelihoods of Somalia that have been affected by erratic rainfall and Desert Locust, the 2020 *Deyr* season cereal harvest and seasonal agricultural employment and incomes have been affected significantly, resulting in moderate to large food consumption gaps both during the current and projection periods.
- In riverine livelihoods of southern Somalia which have been affected by continued flooding both during the 2020 *Gu* and the 2020 *Deyr* seasons, farmers have experienced significant declines in crop harvest and income from agricultural employment. As a result, a significant proportion of poor households in riverine livelihoods face moderate to large food consumption gaps both during the current and projection periods.
- Poor performance of the 2020 Deyr rain season rainfall and its adverse impact on rural livelihoods has also negatively impacted food security among Internally displaced persons (IDPs) in rural areas. Further more, a majority of IDPs in urban areas are poor and, with limited livelihood assets and options and greater reliance on external humanitarian assistance. As a result, a significant proportion of IDPs (both rural and urban) face moderate to large food consumption gaps. Some of the urban poor across Somalia continue to face moderate to large food consumption gaps. For both groups (i.e. urban IDPs and urban poor), the socio-economic impact of COVID-19 was an additional factor (decline in remittances, increased food prices, decline in employment and income earning opportunities in urban areas).


# Food Security Outcomes and Projections (Jul 2020-Jun 2021)



- Based on results of the 2020 Post Deyr assessments, the estimated number of people in urgent need is currently **1.6 million** (Jan-Mar 2021), **even in the presence of humanitarian assistance**. This number would have been higher if there has not been sustained and large scale food assistance.
- Considering various risk factors (likely below average 2021 Gu season rainfall, continued Desert Locust infestation and socio-economic impacts of COVID-19), this number is expected to increase to **2.65 million** (Apr-Jun 2021).

# Humanitarian Assistance and Government Support

**Food Assistance Coverage: July to December 2020**  
(FSC Data)



- Food assistance reached **1.6 million to 1.9 million** people between July and December or an average of **1.8 million** people per month.
- Government/WFP safety net in urban areas (Banadir) is reaching **125 000 people** every month since July 2018 (\$35/month/HH). Government safety net in rural areas (Baxnano/resilience, reached **440 900 people** between July to December 2020 (\$20/month/HH) .
- Life-saving curative and preventive services have been implemented at scale throughout 2020.
- Large scale and sustained humanitarian assistance and government support have contributed to preventing the worsening of food security and nutrition outcomes across many parts of Somalia.




## Nutrition Cluster Data: Life-saving preventive nutrition services for vulnerable population (Jan-Dec 2020)

Number of boys and girls (06-23 months) receiving preventive services through supplementary nutrition products	736 000
Number of pregnant and lactating women and girls (15-49 Years) receiving preventive services through supplementary nutrition products	317 500
Number of pregnant and lactating women receiving first individual counselling on appropriate IYCF/IYCF-E	1 039 600

## Nutrition Cluster Data: Life-saving curative nutrition services (Jan-Dec 2020)

Number of boys and girls 6-59 months with Severe Acute Malnutrition	252 800
Number of boys and girls 6-59 months with Moderate Acute Malnutrition	481 700
Number of PLW with Moderate Acute Malnutrition Treated	144 700

# Food Security Outcomes: Recent Trends Among Rural, Urban and IDP Populations



# Key Messages

- In conclusion, levels of acute food insecurity and malnutrition in Somalia remain high.
- Large scale and sustained humanitarian assistance and government support are preventing worse food security and nutrition outcomes.
- An estimated **838 900** children under the age of five years (total acute malnutrition burden) face acute malnutrition over the next 12 months (Jan-Dec 2021), including **143 200** likely to be severely malnourished. Urgent nutrition and health support is required to address their needs. Urgent health and nutrition support is also required for areas with high prevalence of acute maternal malnutrition.
- Coverage of health and nutrition services (both treatment and prevention), including vitamin A supplementation and measles vaccination should be expanded, especially in areas where coverage is currently low.
- Currently (January-March 2021), an estimated **1.6 million** people are classified in Crisis (IPC Phase 3) and Emergency (IPC Phase 4) and need urgent humanitarian assistance. This number is expected to increase to **2.65 million** people in Crisis (IPC Phase 3) and Emergency (IPC Phase 4) between April and June 2021 due to multiple risk factors.
- Ongoing humanitarian assistance must be scaled up and sustained through mid-2021 to address urgent needs. Livelihoods support is also required for people that are likely to be Stressed or worse (IPC Phase 2 or higher). Population in Stressed (IPC Phase 2) could slide into Crisis or Emergency when they are unable to cope with shocks.
- Drought conditions are developing in parts of northern and southern regions and these are expected to get worse through the January to March 2021 *Jilaal* season. With likely below average Gu season rainfall, drought conditions could worsen.

---

# Thank you

**Somalia IPC Core Team Members: FSNAU/FAO, FEWS NET, WFP/VAM, Food Security Cluster**



Food and Agriculture  
Organization of the  
United Nations



**FSNAU**  
Food Security and Nutrition  
Analysis Unit - Somalia

*Information for Better Livelihoods*



**FEWS NET**  
FAMINE EARLY WARNING SYSTEMS NETWORK



**vam**  
food security analysis



**SOMALIA**  
**FOOD SECURITY CLUSTER**  
*Strengthening Humanitarian Response*